arXiv:0712.1493v1 [nucl-th] 10 Dec 2007

Few-Body Systems 0, 1–3 (2008)

Few-
Body
Systems

© by Springer-Verlag 2008
Printed in Austria

Longitudinal response function of ^4He with a realistic force

S. Bacca [1] *, N. Barnea [2] , W. Leidemann [3], G. Orlandini [3]

[1] Gesellschaft für Schwerionenforschung, Planckstr. 1, 64291 Darmstadt, Germany
[2] Racah Institute of Physics, Hebrew University, Jerusalem, Israel and Institute of Nuclear Physics, University of Washington, Seattle, WA 98195, USA
[3] Dipartimento di Fisica, Università di Trento and INFN (Gruppo Collegato di Trento), Italy

Abstract. The longitudinal response function of ^4He is calculated with the Argonne V18 potential. The comparison with experiment suggests the need of a three-body force. When adding the Urbana IX three-body potential in the calculation of the lower longitudinal multipoles, the total strength is suppressed in the quasi-elastic peak, towards the trend of the experimental data.

The inclusive electron scattering is governed by two kinds of response functions: the longitudinal, $R_L(\omega, \mathbf{q})$, and the transverse response, $R_T(\omega, \mathbf{q})$, where ω and $\mathbf{q}$ are the energy and momentum transfer. $R_{L/T}$ are induced by the electromagnetic charge, $\hat{\rho}(\mathbf{q})$, and current, $\hat{\mathbf{J}}(\mathbf{q})$, operators, respectively. We investigate the medium-low momentum transfer region of the longitudinal response function of ^4He starting with a realistic two-body interaction only, the Argonne V18 (AV18) potential [1], and then adding the Urbana IX (UIX) three-body force [2] as a second step. The longitudinal response is in fact a rich soil for the investigation of three-nucleon force effects, since one can trust the non-relativistic limit and one does not need to include meson exchange currents, as required for R_T to ensure gauge invariance. The longitudinal response is defined as

$$R_L(\omega, \mathbf{q}) = \sum\!\!\!\!\!\!\int_f \left| \langle \Psi_f | \, \hat{\rho}(\mathbf{q}) \, | \Psi_0 \rangle \right|^2 \delta \left(E_f - E_0 - \omega + \frac{\mathbf{q}^2}{2M} \right) , \qquad (1)$$

where $E_{0/f}$ are the energies of the ground and final states $\left| \Psi_{0/f} \right\rangle$ and M is the mass of ^4He. In our calculation of R_L the final state interaction of the continuum four-body wave function is fully taken into account via the Lorentz Integral Transform (LIT) method [3], where one does not calculate (1) directly, but one first evaluates an integral transform of it. The method leads to the solution of a Schrödinger-like equation with bound state-like asymptotics. We solve it making use of a spectral resolution method based on the construction of an effective

---

* *E-mail address:* bacca@gsi.de



Figure 1. Percentage contributions of multipoles of order J to the total strength.

interaction in the hyperspherical harmonics basis (EIHH) [4, 5]. The response function itself is then obtained by a stable inversion of the integral transform.

In the present calculation, as done in [6], we separate the charge operator into isoscalar and isovector parts, which we expand in multipoles. In Fig. 1 we show the total strength of each multipole for two different momentum transfer values, q = 300 MeV/c and q = 500 MeV/c. One notes that the convergence of the multipole expansion gets slower with increasing momentum transfer. However, with the present approach we can easily access the full response function of each multipole as a function of the energy, as presented in Fig. 2 for the same



Figure 2. Isoscalar and isovector response functions of ^4He for different multipoles, consecutively summed up to the indicated order J, for q = 300 MeV/c and q = 500 MeV/c.

two momentum transfer values. Summing then the total isoscalar and isovector contributions, we can directly compare our calculation with the available experimental data. In Fig. 3 one can note that the AV18 two-body potential alone leads to an overprediction of the quasi-elastic peak, especially for q = 300 MeV/c, suggesting the need of a three-nucleon force. By including the UIX three-body force in the calculation of only the monopole ($C_0^{S/V}$) and dipole (C_1^V) responses for q = 300 MeV, in fact we observe a reduction of the total strength. Since these lower multipoles, where the effect of UIX is taken into account, contribute to about 50% of the total strength (see Fig. 1), we can estimate the global effect of the UIX force to be about twice the effect shown in Fig. 3. In future it remains to be seen whether the AV18+UIX nuclear force model is able to reproduce electron scattering experimental data for the longitudinal response in the whole quasi-elastic peak region and for different momentum transfer values.



Figure 3. Longitudinal response function of ${}^4He$ at momentum transfer q = 300 MeV/c and q = 500 MeV/c with the AV18 potential in comparison with experimental data from Bates [7] and Saclay [8]. The curve denoted with AV18+UIX includes the three-body force only in the $C_0^{S/V}$ and C_1^V multipoles.

References

1. Wiringa, R. B., Stoks, V. G. J., Schiavilla, R.: Phys. Rev. **C51**, 38 (1995)
2. Pudliner, B. S., et al.: Phys. Rev. **C56**, 1720 (1997)
3. Efros, V. D., Leidemann, W., Orlandini, G.: Phys. Lett. **B338**, 130 (1994)
4. Barnea, N., Leidemann, W., Orlandini, G.: Phys. Rev. **C61**, 054001 (2000)
5. Barnea, N., Novoselsky, A.: Ann. Phys. (N.Y.) **256**, 192 (1997)
6. Bacca, S., Arenhövel, H. , Barnea, N., Leidemann, W., Orlandini, G.: Phys. Rev. **C76**, 014003 (2007)
7. Dytman, S. A., et al.: Phys. Rev. **C38**, 800 (1988)
8. Zghiche, A., et al.: Nucl. Phys. **A572**, 513 (1994)